

NOTE

Additions and Amendments to the Benthic Marine Algal Flora of Fiji, Including the Island of Rotuma

G. ROBIN SOUTH

ANTOINE DE RAMON N'YEURT and

RAVI ASHNI RAJ-PRASAD

*Marine Studies Programme, The University of the South Pacific
P.O. Box 1168, Suva, Republic of Fiji*

Abstract—Fifty-two new records of benthic marine algae from the Fiji Islands (including the first algal records from the island of Rotuma) are reported, comprising 16 Cyanophyceae, 8 Chlorophyceae, 5 Phaeophyceae, and 23 Rhodophyceae. Included among the new records are species found for the first time at sites other than the type locality (*Streblonema minutulum* Heydrich described from Papua New Guinea, and *Meristotheca procumbens* P. Gabrielson et Kraft, described from Lord Howe Island), and others showing a disjunct northern–southern hemisphere distribution [e.g. *Phaeophila dendroides* (P. Crouan et H. Crouan) Battlers, and *Rosenvingea orientalis* (J. Agardh) Børgesen]. With this list and previously published reports, a total of 383 species of benthic algae is now known from Fiji; on-going studies indicate that this might represent 65–70% of the flora; the cyanophycean flora is especially rich and as yet inadequately described.

Introduction

South & Kasahara (1992) listed a total of 314 benthic marine algae recorded from the Fiji Islands; since then an additional ten taxa have been reported (South, 1991; 1993; South & N'Yeurt, 1993), bringing the total to 324. Recent studies from the Great Astrolabe Reef (Littler & Littler, pers. comm.), Rotuma (N'Yeurt, 1993) and other localities in Fiji, as well as a study of mangrove-associated algae on Viti Levu (Raj, 1993) have resulted in numerous new records, indicating that the flora is much richer than earlier reports suggested.

In this paper we report fifty-two new records of benthic marine algae from Fiji (including Rotuma), and an additional seven taxa listed in Garbary et al. (1991) and omitted from South & Kasahara (1992). Full details of the records from the Island of Rotuma are given in N'Yeurt (1993); these represent the first collections from this isolated location. Raj (1993) reports in detail on the mangrove-associated species.

Materials and Methods

Collecting sites for species listed here are shown in Figure 1. Specimens were obtained with the aid of SCUBA, or by snorkeling, reef walking or examination of scrapings from mangrove pneumatophores. Specimens were preserved in 3–5% formaldehyde in sea water and returned to the laboratory for detailed examination. Voucher specimens are retained as microscope slides, herbarium specimens or liquid-preserved specimens and are housed in the Phycological Herbarium, South Pacific Regional Herbarium, Suva, Fiji (USP). Camera lucida drawings were made with the aid of an Abbé Drawing Tube (Carl Zeiss) and photomicrographs were taken with a Zeiss Photoscope III.

Results

SYSTEMATIC LIST

The arrangement of the list follows that of Silva et al. (1987) where appropriate, and is the same as that adopted in South & Kasahara (1992). In addition to new records, seven species reported by Garbary et al. (1991) and omitted from




Figure 1. Map of the Fiji Islands, with Rotuma (Inset), showing locations where samples were collected. 1. Yageta Island; 2. Ba River; 3. Toberua Passage; 4. Rewa River; 5. Suva Lagoon; 6. Nukulau Island; 7. Makuluva Island; 8. Nukulau Channel; 9. Maka Bay; 10. Ahau; 11. Jolmea; 12. Ropure; 13. Hafea; 14. Lopta; 15. Oinafa; 16. Tuakoi; 17. Hapmafau; 18. Kilinga; 19. Fapufa.

South & Kasahara (1992) are also included. These omissions were a result of changes in Garbary et al. (1991) following revision of their manuscript, and not seen by South & Kasahara (1992) until after the publication of both lists. Records from Fiji proper, including those of Raj (1993) are described here in full. Those from Rotuma are limited to mention of representative material and information on distribution: descriptions and illustrations are given in N'Yeurt (1993) and will be published elsewhere. Accession numbers to specimens housed in the Phycological Herbarium, South Pacific Regional Herbarium are given wherever possible; numbers beginning with "S" refer to slide collections.

CYANOPHYCEAE

CHROOCOCCALES

Chroococcaceae

Aphanocapsa Nägeli

Aphanocapsa sp.

REPRESENTATIVE MATERIAL: Ahau, Rotuma [*N'Yeurt*, 10th Feb., 1993: USP S3:12].

DISTRIBUTION: Cosmopolitan (Bourrelly, 1970).

Gomphosphaeria Kützing

Gomphosphaeria sp.

REPRESENTATIVE MATERIAL: Ahau, Rotuma [*N'Yeurt*, 10th Feb., 1993: USP S3:12].

DISTRIBUTION: Cosmopolitan (Bourrelly, 1970).

Entophysalidaceae

Entophysalis Kützing

Entophysalis sp.

REPRESENTATIVE MATERIAL: Ahau, Rotuma [*N'Yeurt*, 10th Feb., 1993: USP S3:12].

DISTRIBUTION: Cosmopolitan (Bourrelly, 1970).

PLEUROCAPSALES

Hyellaceae

Dermocarpa P. Crouan et H. Crouan

Dermocarpa sp.

REPRESENTATIVE MATERIAL: Ahau, Rotuma [*N'Yeurt*, 10th Feb., 1993: USP S3:12].

DISTRIBUTION: Cosmopolitan (Bourrelly, 1970).

Oncobrysa Meneghini

[= *Hydrococcus* sensu Bourrelly, 1970)

Oncobrysa sp.

REPRESENTATIVE MATERIAL: Ahau, Rotuma [*N'Yeurt*, 10th Feb., 1993: USP S3:12].

DISTRIBUTION: Cosmopolitan (Bourrelly, 1970).

Pleurocapsa Thuret ex Hauck

Pleurocapsa sp.

REPRESENTATIVE MATERIAL: Ahau, Rotuma [*N'Yeurt*, 10th Feb, 1993: USP S3:12].

DISTRIBUTION: Cosmopolitan (Bourrelly, 1970).

CHAMAESIPHONALES

Chamaesiphonaceae

Chamaesiphon A. Braun et Grunow

Chamaesiphon sp.

REPRESENTATIVE MATERIAL: Lopta, Rotuma [*N'Yeurt*, 18th Mar., 1992: USP S4:1].

DISTRIBUTION: Cosmopolitan (Bourrelly, 1970).

Siphononemataceae

Siphononema Geitler

S. polonicum Geitler

REPRESENTATIVE MATERIAL: Lopta, Rotuma [*N'Yeurt*, 18th Mar., 1992: USP S4:1].

DISTRIBUTION: Europe (Bourrelly, 1970:342, pl. 89 figs. 14–16).

NOSTOCALES

Nostocaceae

Anabaena Bory de St. Vincent

Anabaena sp.

REPRESENTATIVE MATERIAL: Ahau, Rotuma [*N'Yeurt*, 10th Feb, 1993: USP S3:12].

DISTRIBUTION: Cosmopolitan (Bourrelly, 1970).

Oscillatoriaceae

Lyngbya C. Agardh

L. aestuarii (Mertens) Liebermann

REPRESENTATIVE MATERIAL: Hapmafau, Rotuma [*N'Yeurt*, 10th Mar., 1992: USP S4:2].

DISTRIBUTION: Cosmopolitan (Lindstedt 1943: 82, pl. 10 figs. 3–4).

Oscillatoria Vaucher

Oscillatoria sp

REPRESENTATIVE MATERIAL: Ahau, Rotuma [*N'Yeurt*, 10th Feb, 1993: USP S3:12].

DISTRIBUTION: Cosmopolitan (Bourrelly, 1970).

Schizothrix Kützing

Schizothrix sp.

REPRESENTATIVE MATERIAL: Hapmafau, Rotuma [*N'Yeurt*, 10th Mar., 1992: USP S3:16].

DISTRIBUTION: Cosmopolitan (Bourrelly, 1970).

Spirulina Turpin

S. tenerima Kützing

REPRESENTATIVE MATERIAL: Hafea, Rotuma [*N'Yeurt*, 5th Mar., 1992: USP S3:17].

DISTRIBUTION: Cosmopolitan (Lindstedt 1943: 55, pl. 6 figs. 12–13).

S. subtilissima Kützing

REPRESENTATIVE MATERIAL: Ropure, Rotuma [*N'Yeurt*, 28th Jan., 1993: USP S3:14].

DISTRIBUTION: Cosmopolitan (Lindstedt 1943: 55, pl. 6 figs. 10–11; Dawson 1957: 128, fig. 311; Umezaki 1961: 65, pl. 10 fig. 7; Tsuda and Wray 1977: 92; Payri and Meinesz 1985a: 504).

Rivulariaceae

Homeothrix (Thuret) Kirchner

Homeothrix sp.

REPRESENTATIVE MATERIAL: Lopta, Rotuma [*N'Yeurt*, 18th Mar., 1992: USP S4:1].

DISTRIBUTION: Cosmopolitan (Bourrelly, 1970:).

Scytonmenataceae

Plectonema Thuret

Plectonema sp.

REPRESENTATIVE MATERIAL: Tuakoi, Rotuma [*N'Yeurt*, 26th Dec., 1992: USP S3:13].

DISTRIBUTION: Cosmopolitan (Bourrelly, 1970:).

Scytonema C. Agardh

Scytonema hofman-bangii C. Agardh

[Garbarry et al., 1991]

CHLOROPHYCEAE

CTENOCLADALES

Ulvellaceae

Acrochaete N. Pringsheim

A. viridis (Reinke) R. Nielsen

[Garbarry et al., 1991]

Phaeophila Hauck

P. dendroides (P. Crouan et H. Crouan) Batters

Figure 2.

REPRESENTATIVE MATERIAL: Bega Lagoon (ex Suva Municipal Market) [South, 12th Oct., 1992; USP S1:9].

Endophytic in the thallus of *Hypnea pannosa* J. Agardh, a species regularly collected for human consumption (South, 1993). Filaments penetrate the inter-cellular spaces of the host, the uppermost lying close to or at the host surface and producing numerous unsheathed hairs from the dorsal surface. Chloroplast parietal, cells containing 1–2 pyrenoids.

DISTRIBUTION: Beqa, Fiji; Northeastern N. America and eastern tropical N. America (Taylor, 1957; 1960); Seychelles Islands (Cöetivy Is.; Kalogina-Gutnik et al., 1992: 17).




Figure 2. *Phaeophila dendroides* endophytic in *Hypnea pannosa*. a. Viewed from the surface. b. Cross section of host showing filaments, with hair cells positioned between epidermal cells and emerging above the host surface. Scales = 50 μm .

CLADOPHORALES Cladophoraceae

Cladophora Kützing

C. conferta P. Crouan et H. Crouan

REPRESENTATIVE MATERIAL: Jolmea, Rotuma [*N'Yeurt*, 21st Jan., 1993: USP 440].

DISTRIBUTION: Tropical Atlantic (van den Hoek 1982: 173, figs. 332-354); Philippines (Silva et al. 1987: 97); Rotuma.

Rhizoclonium Kützing

R. africanum Kützing

REPRESENTATIVE MATERIAL: Hapmafau, Rotuma [*N'Yeurt*, 3rd Feb., 1993: USP 448].

DISTRIBUTION: Solomon Islands (Womersley & Bailey 1970: 265); Northern Australia (Lewis 1987: 15); Philippines (Silva et al. 1987: 99); Rotuma.

R. grande Børgesen

REPRESENTATIVE MATERIAL: Hapmafau, Rotuma [*N'Yeurt*, 17th Mar., 1992: USP S4:5].

DISTRIBUTION: India (Børgesen 1935: 14, figs. 5-6); Tanzania (Jaasund 1976: 5, fig. 12); Philippines (Silva et al. 1987: 99); Japan (Tsuda 1991: 43); Rotuma.

SIPHONOCLADALES

Siphonocladaceae

Cladophoropsis Børgesen

C. sundanensis Reinbold

REPRESENTATIVE MATERIAL: Ba and Rewa Delta, Viti Levu (Raj, 1993: USP S3:7); Fapufa, Rotuma [*N'Yeurt*, 12 Dec., 1991: USP S447; 21 Apr., 1992: USP S4:6].

DISTRIBUTION: Solomon Islands (Womersley & Bailey 1970: 268); Tanzania (Jaasund 1976: 11, fig. 24); Micronesia (Tsuda & Wray 1977: 96); China (Meiling & Tseng 1984: 274, pl. 136 fig. 1); Tahiti (Payri & Meinesz 1985a: 509); Australia (Lewis 1987: 10 ["*sudanensis*"]); Philippines (Silva et al. 1987: 101); Rotuma.

BRYOPSIDALES

Bryopsidaceae

Bryopsis Lamouroux

B. plumosa (Hudson) C. Agardh

REPRESENTATIVE MATERIAL: Hapmafau, Rotuma [*N'Yeurt*, 22nd May, 1992: USP 337; USP 338].

DISTRIBUTION: Lord Howe Island (Lucas 1935: 198); Tropical America (Taylor 1960: 131, pl. 9 fig. 11); Micronesia (Tsuda & Wray 1977: 94); Philippines (Silva et al. 1987: 103); New Caledonia (Garrigue & Tsuda 1988: 57); Rotuma.

Halimiaceae

Halimeda Lamouroux

H. micronesica Yamada

REPRESENTATIVE MATERIAL: Oinafa, Rotuma [*N'Yeurt*, 19th Mar., 1992: USP 394, 396; 17th May, 1992: USP 395].

DISTRIBUTION: Solomon Islands (Womersley & Bailey 1970: 282); Micronesia (Tsuda & Wray 1977: 98); China (Meiling & Tseng 1984: 290, pl. 144, fig. 1); Polynesia (Payri & Meinesz 1985b: 643, figs. 16, 18, 24, 46); Rotuma.

Udoteaceae

Chlorodesmis Harvey ex Bailey

C. major Zanardini

REPRESENTATIVE MATERIAL: Hapmafau, Rotuma [*N'Yeurt*, 10th Mar., 1992: USP 340].

DISTRIBUTION: Lord Howe Island (Lucas 1935: 200); Hawaii (Egerod 1952: 377, fig. 9c); Northern Australia (Lewis 1987: 27); Philippines (Silva et al. 1987: 118); Rotuma.

Rhipiliopsis A. Gepp et E. S. Gepp

Rhipiliopsis sp.

[Garbary et al., 1991]

DASYCLADALES

Polyphysaceae

Acetabularia Lamouroux*A. parvula* Solms-Laubach[Garbary et al., 1991 (as *A. moebii* Solms-Laubach)]

PHAEOPHYCEAE

ECTOCARPALES

Ectocarpaceae

Feldmannia Hamel*F. columellaris* (Børgesen) Islam


[Garbary et al., 1991]

Hincksi J. E. Gray*H. breviarticulata* (J. Agardh) P. C. Silva[=*Ectocarpus breviarticulata* J. Agardh; see Silva et al. 1987: 73].REPRESENTATIVE MATERIAL: Lopta, Rotuma [*N'Yeurt*, 18th Mar., 1992: USP 357].

DISTRIBUTION: Philippines (Silva et al. 1987: 73); Hawaii (Abbott 1989: 226); Solomon Islands (Womersley & Bailey 1970: 288); Micronesia (Tsuda & Wray 1977: 101); China (Baoren & Tseng 1984: 168, pl. 85 fig. 1); Northern Australia (Lewis 1985: 3); Polynesia (Payri & Meinesz 1985: 505); Nauru (South & Yen 1992: 128); Rotuma.

Streblonema Derbès et Solier in Castagne*S. minutulum* Heydrich

Fig. 3.

Heydrich, 1892: 471, pl. 26, fig. 15; Kuckuck, 1954: 110–111, fig. 5. [type locality: Papua New Guinea, endophytic in *Sebdenia ceylandica*]Figure 3. *Streblonema minutulum* endophytic in the outer layers of *Soliera robusta*.

Narrow hypha-like filaments penetrate the host intercellular spaces; emergent are assimilators (a), phaeophycean hairs (p) and plurilocular sporangia (pl).

Scale = 50 μ m.

REPRESENTATIVE MATERIAL: Suva Lagoon, Viti Levu (ex Suva Municipal Market) [South, 12th Oct., 1992; USP S1:8]. Endophytic in *Solieria robusta* (Greville) Kylin. Specimens bore plurilocular sporangia, phaeophycean hairs and short vegetative filaments of 3–4 cells emergent above the host surface. Endophytic basal filaments more or less colourless, narrow, frequently branched and forming a network between the host cortical cells. Plants closely resemble those illustrated in Kuckuck (1954, fig. 5).

DISTRIBUTION: Papua New Guinea; Fiji.

DICTYOTALES

Dictyotopsidae

Dictyotopsis Troll

D. propagulifera Troll

Troll, 1931: 474, figs. 4–18 [type locality Island of Belawan, Indonesia]; Post, 1967: 279; Farrant & King, 1989: 369; King, 1990: 58.

REPRESENTATIVE MATERIAL: Ba, Rewa Delta, Viti Levu [Raj, 1992; USP S2:17]

Occurs in the upper, drier region of the mangal, and was present on all mangrove species examined in the Rewa River Delta, Viti Levu, but only on *Rhizophora samoensis* (Hochr.) Salvoza and *R. stylosa* Griff. pneumatophores in the Ba River, Viti Levu (Raj, 1993: 64). The single apical cells segment laterally, and the thallus is monostromatic.

DISTRIBUTION: Indonesia; Fiji (first recorded in Post, 1967); Australia (New South Wales; Farrant and King, 1989); Fiji.

DICTYOSIPHONALES

Chnoosporaceae

Chnoospora J. Agardh

C. minima (Hering) Papenfuss

REPRESENTATIVE MATERIAL: Lopta, Rotuma [N'Yeurt, 18th Mar., 1992; USP 358; 15 Dec., 1992; USP 478, 479].

DISTRIBUTION: Solomon Islands (Womersley & Bailey 1970: 293); Hawaii (Abbott 1989: 227); China (Baoren & Tseng 1984: 184, pl. 93 fig. 2; Northern Australia (Lewis 1985: 8); Rotuma.

SCYTOSIPHONALES

Scytosiphonaceae

Rosenvingea Børgesen

R. orientalis (J. Agardh) Børgesen

Fig. 4.

Børgesen, 1914: 26.

Asperococcus orientalis J. Agardh, 1848: 78 [type locality Manila, Luzon, Philippines]. For nomenclature and synonymy, see Silva et al. (1987: 80).

REPRESENTATIVE MATERIAL: Known from only a single specimen in Fiji, from Nukulau Is., Suva Lagoon [South, 9 Sept., 1991; USP 232].

In a silty channel between Nukulau and Makaluva Islands, Suva Lagoon, Viti Levu, just below low tide level and attached by small discoid holdfasts to a fragment of dead *Acropora* coral. Plants to 18cm tall, branching frequent, dichotomous to subdichotomous, terminating in fine, sharply attenuated tips. Thallus hollow, phaeophycean hairs numerous; cortex of 1-2 layers of isodiametric-angular cells, epidermis single-layered. Fertile plants not found.

DISTRIBUTION: Philippines (Silva et al., 1987: 80); Eniwetok Atoll (Dawson 1957); Fiji.

RHODOPHYCEAE

Florideophycidae

ACROCHAETIALES

Acrochaetiaceae

Audouinella Børy

A. polyblasta (Rosenvinge) J. Price, Lawson et John

[= *Acrochaetium sargassi* Børgesen, vide Woelkerling 1973: 84]

REPRESENTATIVE MATERIAL: Hapmafau, Rotuma [*N'Yeurt*, 10th Mar., 1992: USP S4:7].

DISTRIBUTION: North America (South & Tittley 1986:3) Indonesia (Weber-van Bosse 1921: 193, as *Acrochaetium sargassi* Børgesen); Solomon Islands (Womersley & Bailey 1970: 301, as *Acrochaetium sargassi* Børgesen); Rotuma.

A. saviana (Meneghini) Woelkerling

Woelkerling, 1973: 560, figs. 56-60.

For nomenclature and synonymy, see Woelkerling (1973).

REPRESENTATIVE MATERIAL: Suva Lagoon [South, 8th April, 1993: USP568].

An abundant epiphyte on the leaves of the seagrass *Halodule uninervis* (Forsskål) Ascherson and on the filaments of *Chaetomorpha crassa* (C. Agardh) Kützing. Plants to 500 µm tall, the original spore non-persistent; prostrate system as a limited, pseudo-parenchymatous disk; erect filaments freely, irregularly branched, unicellular hairs lacking; chloroplast single, parietal and one pyrenoid present; monosporangia frequent, single, in secund series along the laterals, sessile or stalked.

DISTRIBUTION: Nearly cosmopolitan (Woelkerling 1973: 561).

GELIDIALES

Gelidiaceae

Gelidium Lamouroux

G. crinale (Turner) Gaillon

var. *perpusillum* Piccone et Grunow

Piccone, 1884: 317 [type locality: Massawa, Ethiopia]

REPRESENTATIVE MATERIAL: Ba and Rewa Rivers, Viti Levu [*Raj*,




Figure 4. *Rosenvingea orientalis*, showing habit (a), an apical region with phaeophycean hairs (b) and a cross section showing the hollow thallus (c). Scales (a) = 2.0 cm, (b), (c) = 0.5 mm.

1993; S2:12].

Found on the pneumatophores of all mangrove species examined (Raj, 1993: 56).

DISTRIBUTION: A general tropical distribution; Indonesia (Tanaka & Chihara, 1988a, b); Fiji.

CORALLINALES

Corallinaceae

Jania Lamouroux

J. rubens (Linnaeus) Lamouroux

REPRESENTATIVE MATERIAL: Lopta, Rotuma [N'Yeurt, 15 Dec., 1992: USP 559].

DISTRIBUTION: Solomon Islands (Womersley & Bailey 1970: 314); Greece (Haritonidis & Tsekos 1976: 281); Micronesia (Tsuda & Wray 1977: 108);

Northern Australia (Lewis 1984: 15); Polynesia (Payri & Meinesz 1985: 513); Philippines (Silva et al. 1987: 35); Rotuma.

Mesopyhyllum Lemoine

M. mesomorphum (Foslie) Adey

REPRESENTATIVE MATERIAL: Hapmafau, Rotuma [*N'Yeurt*, 10th Mar., 1992: USP 450]; Kilinga, Rotuma [*N'Yeurt*, 22nd Dec., 1992: USP 440].

DISTRIBUTION: Guam (Gordon et al. 1976: 252, pl. II figs. 1-3); Micronesia (Tsuda & Wray 1977: 110); Hawaii (Magruder & Hunt 1979: 85); Northern Australia (Lewis 1984: 17); Rotuma.

GIGARTINALES

Plocamiaceae

Plocamium Lamouroux

P. cartilagineum (Linnaeus) Dixon

Fig. 5.

Dixon, 1967: 58.

Fucus cartilagineus Linnaeus, 1753: 1161 [type locality: northern Europe]. For nomenclature, synonymy and description, see South and Adams, 1979: 124, figs. 2, 11-15.

REPRESENTATIVE MATERIAL: Yageta Island [*Würtz*, 19th Sept., 1992; USP 567]; Nukulau Channel, Viti Levu [*Carlson*, 20 Aug., 1972; USP 314]; Toberua Passage, Viti Levu [*Carlson*, 3 Dec., 1972; USP 316]; Makuluva Island, Viti Levu [*South*, 9 Sept., 1991; USP 315].

Plants closely fit the description of this species in South & Adams (1979: 124); ramuli are in alternate series of 3-4, and adventitious ramuli opposite to or between members of the alternating series are absent. The lowermost ramulus is simple, and not hooked in form. Tetrasporangial stichidia are pedicillate in the axils of ramuli, or grouped on the upper margins of ramuli and along the axis; they are at first simple, becoming branched and elongate.

DISTRIBUTION: According to Womersley (1971) this is the only very widespread species of *Plocamium*, being recorded from many temperate seas; its presence in Fiji is thus of interest. Occurs throughout New Zealand and extends as far south as Antarctica (South & Adams, 1979: 125); also occurs in Europe, N. America and Australia (Womersley, 1971: 24). Previous records of *Plocamium* from Fiji have been attributed to *P. abnorme* Hooker fil. et Harvey (South & Kasahara, 1992: 62), a species later placed in synonymy with *P. angustum* (J. Agardh) Hooker fil. et Harvey (South & Adams, 1979: 121). Tetrasporic collections from these sites are needed to re-examine these records in the light of this report of *P. cartilagineum*.


Gracilariaeae

Gracilaria Greville

G. arcuata Zanardini

REPRESENTATIVE MATERIAL: Lopta, Rotuma [*N'Yeurt*, 18th Mar., 1992: USP 349].

DISTRIBUTION: Indonesia (Weber-van Bosse 1928: 429, fig. 173); Tan-



Figures 5–10. Figure 5. *Plocamium cartilagineum* showing elongate, branched tetrasporangial stichidia. Scale = 50 µm. Figure 6. *Dasyphila plumariooides*. Lateral branches (l) are borne on the main axis (m), in alternate series; these may bear secondary laterals of unlimited growth (sl) also alternately arranged, and uniserrate, alternately branched laterals of limited growth (lm) arranged in distichous, opposite pairs. Scale = 50 µm. Figure 7. *Dasyphila plumariooides*. Details of lateral branch with opposite pairs of branches of limited growth; dorsal and ventral branches are suppressed. Scale = 50 µm. Figure 8. *Dasyphila plumariooides*. Apices of ultimate branchlets, showing that initial development of laterals is on the lower side of branches, becoming even towards the base. Scale = 50 µm. Figure 9. *Myriogramme* sp. showing foliose lateral and tetrasporangial sori. Scale = 500 µm. Figure 10. *Myriogramme* sp. showing monostromatic thallus bearing uniserrate rhizoids (small arrows) and multiserrate haptera with terminal rhizoids (large arrows). Scale = 500 µm.

zania (Jaasund 1976: 85, fig. 173); Micronesia (Tsuda & Wray 1977: 106); Northern Australia (Lewis 1984: 24); China (Tseng et al. 1984: 102, pl. 54 fig. 1); Philippines (Silva et al. 1987: 40); New Caledonia (Garrigue & Tsuda 1988: 65); Rotuma.

Solieriaceae

Meristotheca J. Agardh

M. procumbens P. Gabrielson et Kraft

REPRESENTATIVE MATERIAL: Tuakoi, Rotuma [*N'Yeurt*, 2nd Mar., 1992: USP 351].

DISTRIBUTION: Lord Howe Island (Gabrielson & Kraft 1984: 241, fig. 14A-D); Rotuma.

RHODYMENIALES

Rhodymeniaceae

Coelarthrum Børgesen

C. boergesenii Weber-van Bosse

REPRESENTATIVE MATERIAL: Ropure, Rotuma [*N'Yeurt*, 28 Jan., 1993: USP 441].

DISTRIBUTION: Indonesia (Weber-van Bosse 1928: 473, figs. 207-208); Micronesia (Tsuda & Wray 1977: 105); Northern Australia (Lewis 1984: 37); Hawaii (Abbott 1989: 229); Japan (Tsuda 1991: 50); Rotuma.

Rhodymenia Greville

R. divaricata Dawson

REPRESENTATIVE MATERIAL: Maka, Rotuma [*N'Yeurt*, 17th Feb., 1992: USP 352, 353].

DISTRIBUTION: Mexico (Dawson 1941: 141, pl. 23 fig. 31) Micronesia (Tsuda & Wray 1977: 111); Rotuma.

CERAMIALES

Ceramiaceae

Centroceras Kützing

C. apiculatum Yamada

[Garbary et al., 1991]

Ceramiella Børgesen

Ceramiella sp.

REPRESENTATIVE MATERIAL; Hapmafau, Rotuma [*N'Yeurt*, 10th Mar., 1992: USP S4:10, S4:11, S4:12]; Jolmea, Rotuma [*N'Yeurt*, 28th Jan., 1993: USP S4:9].

DISTRIBUTION: Indonesia (Weber-van Bosse 1923: 322); Philippines (Silva et al. 1987: 54); Rotuma.

Ceramium Roth

C. mazatlanense Dawson

[Garbary et al., 1991]

C. zacae Setchell et Gardner

REPRESENTATIVE MATERIAL: Hapmafau, Rotuma [*N'Yeurt*, 10th Mar., 1992: USP S4:8].

DISTRIBUTION: Mexico (Dawson 1950: 134, pl. 2 figs. 27-28); Ewinetok Atoll, Micronesia (Dawson 1957: 8); Philippines (Fortes & Trono 1979: 60, fig. 11; Silva et al. 1987: 56); Rotuma.

***C. codii* (Richards) Feldmann-Mazoyer**

REPRESENTATIVE MATERIAL: Hafea, Rotuma [*N'Yeurt*, 5th May., 1992: USP S4:14].

DISTRIBUTION: Bermuda (Richards 1901:264, pls. 21–22); Tanzania (Jaasund 1970b: 68, fig. 1, F, N; 1976: 107, fig. 216); Australia (Cribb 1983: 80, pl. 27 figs. 1–4; Lewis 1984: 41; Millar 1990: 393, figs. 41D–F; 43B; Price & Scott 1992: 86, fig. 26A–D); Rotuma.

***Dasyphila* Sonder**

Figs. 6–8.

***D. plumariooides* Yendo**

Yendo, 1920: 7 [type locality Kotosho (Hung-t'ou), Taiwan]. Okamura, 1923: 198, Figs. 5–11; 1931: 52, pl. 277, fig. 11; Womersley and Bailey, 1970: 326, fig. 9; Silva et al., 1987: 56; Lewis and Norris, 1987: 22.

REPRESENTATIVE MATERIAL Yageta Island [*Würtz*, 17th Sept., 1992; USP 566].

Plants from Fiji exactly fit the description in Womersley and Bailey (1970) and the figures in Okamura (1923, 1931). To date, only sterile plants have been found, in a single collection.

DISTRIBUTION: Taiwan, Philippines, Solomon Islands, Fiji.

Delesseriaceae

***Caloglossa* (Harvey) J. Agardh**

See recent discussion of *Caloglossa* in Sheath et al. (1993: 113).

***Caloglossa* sp. 1 inedit**

This species resembles *Caloglossa adnata* (Zanardini) De Toni previously recorded from Fiji (Chapman, 1971; South & Kasahara, 1992: 64). As stated in Raj (1993), however, it differs in the location of the rhizoids, which are formed on any cell of the blade compared with *C. adnata*, where they are produced in groups along the midrib only. King & Puttock (pers. comm.) are in process of describing this as a new species (*C. adhaerens* sp. inedit.) in a forthcoming revision of the genus. The occurrence of *Caloglossa adnata* in Fiji should be reviewed in the light of this.

***Caloglossa* sp. 2 inedit**

This species also resembles *Caloglossa adnata*, but will be described as a new species (*C. bengalensis* sp. inedit.) by King & Puttock in their forthcoming revision of the genus.

Hypoglossum* Kützing**H. caloglossoides* Wynne et Kraft**

REPRESENTATIVE MATERIAL: Tuakoi, Rotuma [*N'Yeurt*, 2nd May., 1992: USP S4:13].

DISTRIBUTION: Lord Howe Island (Wynne and Kraft 1985: 20, figs. 1–19); Australia (Price and Scott 1992: 137, fig. 47A–C); Rotuma.

Myriogramme* Kylin**Myriogramme* sp.**

Figs. 9, 10.

REPRESENTATIVE MATERIAL: Yageta Island [Würtz, 17th Sept. 1992: USP 581; S1:7].

Small plants creeping on the base of *Halimeda*, at a depth of 17 m. Thallus monostromatic, lacking veins or nerves, branching irregular, growth from the margins, diffuse; attachment by multicellular, sometimes robust holdfasts produced from the thallus margins, terminating in multicellular rhizoids; tetrasporangia tetrahedrally divided, in oval or lunate sori on foliose laterals, or in the marginal areas of the main thallus. This may represent an undescribed species.

Rhodomelaceae

Bostrychia Montagne¹

B. simpliciuscula Harvey ex J. Agardh

Agardh 1863: 854 [type locality, Tonga].

REPRESENTATIVE MATERIAL: Ba and Rewa Rivers, Viti Levu [Raj, 1992: USP S2:13]

In Fiji, epiphytic on mangrove pneumatophores, and also occurs as mats on mud in tidal inlets and rivers (Raj 1993). The main axes are polysiphonous and ecorticate, and the ultimate branches are partially monosiphonous. Older prostrate axes have few non layer-forming cortical cells as compared with *B. mortiziana* (Sonder ex Kützing) J. Agardh and *B. radicans* (Montagne) Montagne in Orbigny. The holdfasts are formed from adventitious cortication of pericentral cells (peripherohaptera) and the later branches occur less frequently and are not found on successive axial cells.

DISTRIBUTION: Tropical and sub-tropical coasts of Queensland to 30 °S (King & Puttock 1989); Japan, south-east Asia, oceania (Fiji, Tonga), Papua New Guinea (King 1990) and Kenya (Coppejans & Gallin 1989). Records from Nigeria are misidentifications (King & Puttock 1989).

Chondria C. Agardh

C. sedifolia Harvey

REPRESENTATIVE MATERIAL: Lopta, Rotuma [N'Yeurt, 10th Mar., 1992: USP 475].

DISTRIBUTION: Florida, USA (Dawes 1974: 155); Tanzania (Jaasund 1976: 135, fig. 273a, b: pl. 9); Philippines (Silva et al. 1987: 63); Rotuma.

Laurencia Lamouroux

L. corymbosa J. Agardh

[Garbary et al., 1991]

L. venusta Yamada

REPRESENTATIVE MATERIAL: Lopta, Rotuma [N'Yeurt, 10th Mar., 1992: USP S477].

DISTRIBUTION: Japan (Saito 1967: 14, pls. V and VI; text-figs. 8–14); Tanzania (Jaasund 1970a: 62, fig. 1B; 1976: 141, fig. 284); Australia (Lewis

¹ *Rhodolachne radicans*, described from Fiji by Itono (1985), is now not believed to be representative of the genus *Bostrychia*, and will be re-described (Raj 1993; King, pers. comm.).

1984: 61); Taiwan (Lewis & Norris 1987: 23); Philippines (Silva et al. 1987: 68); Rotuma.

Discussion

The total number of benthic marine algae now reported from Fiji is 383. A significant number of the new records listed here (38 taxa) have resulted from the first detailed phycological survey of the shallow-water algae of Rotuma (N'Yeurt 1993). Situated some 300 km N. of the main Fiji Group, Rotuma experiences a somewhat warmer and less contrasting seasonal climate. In addition, Littler & Littler (pers. comm.) reported that they discovered a substantial number of new records of benthic marine algae from Fiji (including a number of undescribed species) as a result of their 1992 survey of the Great Astrolabe Reef, Kadavu.

The Cyanophyceae listed here require substantial further work; preliminary studies indicate that the cyanophyte flora is very diverse indeed, and many species will be added to those already known from Fiji.

The Fijian marine algal flora includes many species that are characteristic of the Indo-West Pacific region, as well as a number that are pan-tropical or cosmopolitan. However, a number of the species recorded here suggest that our flora also has some unexpected disjunct northern hemisphere-southern hemisphere distributions. Examples are the occurrence of *Phaeophila dendroides* (Chlorophyceae) and *Rosenvingea orientalis* (Phaeophyceae). Other species, to date known only from their type localities, have been reported here (e.g. *Strebblonema minutum* from Papua New Guinea; *Meristotheca procumbens* from Lord Howe Island). *Dictyotopsis propagulifera*, scarcely known before, is now reported to be a widespread species in the Fijian mangal (Raj 1993); the first report of this species from Fiji (Post 1967) was omitted in error by South & Kasahara (1992).

The Fijian flora at present includes no endemics. Several new species have been discovered, however, and are awaiting formal publication (Kasahara 1985, Littler & Littler, unpubl., South et al., unpubl.) and some of these may be endemic.

The rate of discovery of new records continues to be high, and as more surveys (particularly sub-tidal) are carried out many more species may be added to the flora. We estimate that approximately 65–70 percent of the flora is known.

Acknowledgements

All the authors gratefully acknowledge financial support from the University of the South Pacific. ADRN was also supported by an International Centre for Ocean Development (ICOD; Canada) Post Graduate Scholarship; GRS acknowledges the Canadian International Development Agency (CIDA) and ICOD for financial assistance. We thank Dr. B. A. Whitton for his review of the Cyanophyceae, and Dr. G. T. Kraft for examining material of *Meristotheca*. Jürgen Würtz assisted with subtidal collections from the Yasawas and the Great Astrolabe Reef.

References

- Abbott, I. A. 1989. Marine algae of the northwest Hawaiian Islands. *Pacific Sci.* 43: 223–233.
- Agardh, J. G. 1848. *Species genera et ordines algarum. Volumen primum: algas fucoideas complectens.* viii + 363 pp. Lunda [Lund].
- Agardh, J. G. 1863. *Species, genera et ordines algarum. Volumen tertium: genera et ordines floridearum.* Lund.
- Baoren, L. & C. K. Tseng. 1984. *Phaeophyta. In Common Seaweeds of China* (Tseng C.K., ed.), pp. 167–247. Science Press, Beijing.
- Børgesen, F. 1914. The marine algae of the Danish West Indies, Part 2: Phaeophyceae (1): *Dansk. Bot. Arkiv.* 2(2): 1–68, figs. 1–86.
- Børgesen, F. 1935. A List of Marine Algae from Bombay. *Kongelige Danske Videnskabernes Selskab, Biologiske Meddelelser* 12(2): 64 pp., 25 figs, x plates.
- Bourrelly, P. 1970. *Les Algues D'eau Douce. Tome III. Les Algues Bleues et Rouges: Les Eуглéniens, Peridiniens et Cryptomonadines.* Éditions N. Bouée & Cie, Paris. 512 pp.
- Chapman, V. J. 1971. The marine algae of Fiji. *Rev. Algol.* 2: 164–171.
- Coppejans, E. & E. Gallin. 1989. Macroalgae associated with the mangrove vegetation of Gazi Bay (Kenya). *Bull. Soc. Roy. Belg.* 122: 47–60.
- Cribb, A. B. 1983. Marine Algae of the southern Great Barrier Reef. Part 1. Rhodophyta. Australian Coral Reef Society: Brisbane. Handbook No. 2. 173 + [2] pp., 71 pls.
- Dawes, C. J. 1974. Marine Algae of the West Coast of Florida. University of Miami Press. 201 pp.
- Dawson, E. Y. 1941. A review of the genus *Rhodymenia* with descriptions of new species. *Allan Hancock Pacific Expeditions* 3(7/8): 115–181.
- Dawson, E. Y. 1950. A review of *Ceramium* along the Pacific coast of North America with special reference to its Mexican representatives. *Farlowia* 4(1): 113–138.
- Dawson, E. Y. 1957. An annotated list of marine algae from Eniwetok atoll, Marshall Islands. *Pacific Science* 11: 92–132.
- Dixon, P. S. 1967. The typification of *Fucus cartilagineus* L. and *F. corneus* Huds. *Blumea* 15: 55–62.
- Egerod, L. E. 1952. An analysis of the siphonous Chlorophycophyta with special reference to the Siphonocladales, Siphonales and Dasycladales of Hawaii. *Univ. Calif. Pub. Bot.* 25: 325–454.
- Farrant, P. A. & R. J. King. 1989. The Dictyotales (Algae: Phaeophyta) of New South Wales. *Proc. Linn. Soc.* 110: 369–405.
- Fortes, M. D. & G. C. Trono, Jr. 1979. Marine algal microphytes new to the Philippines. *Kalikasan, Philipp. J. Biol.* 8(1): 51–68.
- Gabrielson, P. W. & G. T. Kraft. 1984. The marine algae of Lord Howe Island (N.S.W.): the Family Solieriaceae (Gigartinales, Rhodophyta). *Brunonia* 7: 217–251.

- Garbary, D. J., J. C. Oliveira, R. F. Scagel & S. Villeneuve. 1991. Notes and distribution records for the marine algae of Fiji. *Micronesica* 24: 249–260.
- Garrigue, C. & R. T. Tsuda. 1988. Catalog of marine benthic algae from New Caledonia. *Micronesica* 21: 53–70.
- Gordon, G. D., T. Masaki, & H. Akioka. 1976. Floristic and distributional account of the common crustose coralline algae on Guam. *Micronesica* 12: 247–277.
- Haritonidis, S. & I. Tsekos. 1976. Marine algae of the Greek West coast. *Bot. Mar.* 12: 273–286.
- Heydrich, F. 1892. Ber. D. Bot. Ges. 10: 471.
- Hoek, C. van den. 1982. A Taxonomic Revision of the American Species of *Cladophora* (Chlorophyceae) in the North Atlantic Ocean and Their Geographic Distribution. North-Holland Publishing Company, Amsterdam. 236 pp.
- Itono, H. *Rhodolachne radicosa*, a new species of red alga (Rhodomelaceae, Ceramiales) from Fiji and southern parts of Japan. Kagoshima Univ. Res. Center S. Pac. Occas. Pap. No. 5.
- Jaasund, E. 1970a. Marine algae in Tanzania II. *Bot. Mar.* 13: 59–64.
- Jaasund, E. 1970b. Marine algae in Tanzania II. *Bot. Mar.* 13: 65–70.
- Jaasund, E. 1976. Seaweeds in Tanzania. Univ. Tromsø.
- Kalogina-Gutnik, A. A., L. P. Perestenko & T. V. Titlyanova. 1992. Species composition, distribution and abundance of algae and seagrasses of the Seychelles Islands. *Atoll Research Bull.* 396: 1–67.
- Kasahara, H. 1985. Marine Green and Red Algae Collected at Viti Levu and Ndravuni Islands of Fiji. M. Ag. Thesis, Kyoto University. 83 pp.
- King, R. J. 1990. Macroalgae associated with the mangrove vegetation of Papua New Guinea. *Bot. Mar.* 33: 55–62.
- King, R. J. & C. F. Puttock. 1989. Morphology and taxonomy of *Bostrychia* and *Stictosiphonia* (Rhodomelaceae/Rhodophyta). *Aust. Syst. Bot.* 2: 1–73.
- Kuckuck, P. 1954. Ectocarpaceen Studien II. *Streblonema*. Helgol. Wiss. Meeresunters. 5: 103–117 [pp. 43–57 In Kornmann, P. 1964. Ectocarpaceen Studien von Paul Kuckuck. Biologische Anstalt, Helgoland].
- Lewis, J. A. 1984. Checklist and Bibliography of Benthic Marine Macroalgae recorded from Northern Australia. I. Rhodophyta. Department of Defense: Defense Science and Technology Organisation Materials Research Laboratories Report MRL-R-912. Melbourne. 99 pp.
- Lewis, J. A. 1985. Checklist and Bibliography of Benthic Marine Macroalgae recorded from Northern Australia. II. Phaeophyta. Department of Defense: Defense Science and Technology Organisation Materials Research Laboratories Report MRL-R-962. Melbourne. 40pp.
- Lewis, J. A. 1987. Checklist and Bibliography of Benthic Marine Macroalgae recorded from Northern Australia. III. Chlorophyta. Department of Defense: Defense Science and Technology Organisation Materials Research Laboratories Report MRL-R-1063. Melbourne. 56 pp.

- Lewis, J. E. & J. N. Norris. 1987. A History and Annotated Account of the Benthic Marine Algae of Taiwan. Smithson. Contr. Mar. Sci. 29: 38 pp.
- Lindstedt, A. 1943. Die Flora Der Marinens Cyanophyceen Der Schwedischen Westküste. Håkan Ohlssons Buchdruckerei, Lund. 121 pp. + 11 pls.
- Linnaeus, C. 1753. Species plantarum 2. Stockholm.
- Lucas, A. H. S. 1935. The marine algae of Lord Howe Island. Proc. Linn. Soc. NSW 60: 194–232 + pls 5–9.
- Magruder, W. H. & J. W. Hunt. 1979. Seaweeds of Hawaii. A photographic identification guide. Oriental Publishing Company, Honolulu. 116 pp.
- Meiling, D. & C. K. Tseng. 1984. Chlorophyta. In Common Seaweeds of China (Tseng C. K., ed.), pp. 249–300. Science Press, Beijing.
- Millar, A. J. K. 1990. Marine red algae of the Coffs Harbour region, northern New South Wales. Aust. Syst. Bot. 3: 293–593.
- N'Yeurt, A. D. R. 1993. A Floristic Survey of the Intertidal and Shallow Benthic Marine Algae of Rotuma. M.Sc. Thesis, Univ. S. Pacific. 258 pp.
- Okamura, K. 1923. Icones of Japanese Algae. Vol. IV. 205 pp., plates CLI–CCI. Tokyo.
- Okamura, K. 1931. Icones of Japanese Algae, Vol. VI. 101 + 96 pages, plates CCLI–CCC. Tokyo.
- Payri, C. E. & A. Meinesz. 1985a. Algae. Proc. 5th Int. Coral Reef Symp. 6: 498–518.
- Payri, C. E. & A. Meinesz. 1985b. Taxonomy and distribution of the genus *Halimeda* (Chlorophyta, Caulerpales) in French Polynesia. Proc. 5th Int. Coral Reef Symp. 6: 641–648.
- Piccone, A. 1864. Contribuzioni all'allgologia Eritrea. Nov. Giorn. Bot. Ital. 16: 281–332, plates VII–IX.
- Post, E. 1967. Verbreitung und Ökologie von *Dictyotopsis propagulifera* W. Troll. Rev. Algol. 8: 279–291.
- Price, I. R. & F. J. Scott. 1992. The Turf Algal Flora of the Great Barrier Reef. Part I. Rhodophyta. James Cook University of North Queensland. 266 pp.
- Raj, R. A. 1993. A Floristic Study of Mangrove-associated Algae in Fiji. M.Sc. Thesis, University of the South Pacific. 105 pp. + Appendix.
- Richards, H. M. 1901. *Ceramothamnion codii*, a new rhodophyceous algae. Bull Torrey Bot Club 28: 257–265, pls. 21–22.
- Saito, Y. 1967. Studies on Japanese species of *Laurencia*, with special reference to their comparative morphology. Mem. Fac. Fish. Hokkaido Univ. 15(1): 1–81.
- Setchell, W. A. 1926. Tahitian algae and Tahitian spermatophytes. Univ. Calif. Publ. Bot. 12(5): 61–143.
- Sheath, R. G., M. L. Vis & K. M. Cole. 1993. Distribution and systematics of freshwater Ceramiales (Rhodophyta) in North America. J. Phycol. 29: 108–117.
- Silva, P. C., E. G. Meñez & R. L. Moe. 1987. Catalog of the Marine Benthic Algae of the Philippines. Smithson. Contrib. Mar. Sci. 27: iv + 179 pp.

- South, G. R. 1991. Benthic Marine Algae from Dravuni Island and Astrolabe Islands, Kadavu, Fiji. USP Mar. Stud. Prog. Tech. Rep. 1991(5): 13 pp.
- South, G. R. 1993. Edible seaweeds of Fiji: an ethnobotanical study. Bot. Mar. 36: 335–349.
- South, G. R. and N. M. Adams. 1979. A revision of the genus *Plocamium* Lamouroux (Rhodophyta, Gigartinales) in New Zealand. Phycologia 18(2): 120–132.
- South, G. R. and H. Kasahara. 1992. A preliminary checklist of the benthic marine algae of the Fiji Islands, South Pacific. Micronesica 25: 41–70.
- South, G. R. & A. D. R. N'Yeurt. 1993. Contributions to a catalogue of benthic marine algae of Fiji. II. *Caulerpa* and *Caulerpella* (Chlorophyta–Caulerpales). Micronesica 26: 109–138.
- South, G. R. & I. Tittley. 1986. A checklist and distributional index of the benthic marine algae of the North Atlantic Ocean. Huntsman Mar. Lab. and Br. Mus. (Nat Hist). 76 pp.
- South, G. R. & S. Yen. 1992. Notes on the benthic marine algae of Nauru, Central Pacific. Micronesica 25: 123–131.
- Tanaka, J. & M. Chihara. 1988a. Algal flora associated with mangroves. 1. Species composition and ecology of macroalgae in mangrove brackish areas of east Indonesia. 2. Macroalgal flora in mangrove brackish areas of east Indonesia. In, Ogino, K. & M. Chihara (eds.) Biological System of Mangroves, a Report of East Indonesian Mangrove Expedition, 1986. pp. 21–34. Ehime University, Japan.
- Tanaka, J. & M. Chihara. 1988b. Macroalgae in Indonesian mangrove forests. Bull. Nat. Sci. Mus. Ser. B. 14(3): 93–106.
- Taylor, W. R. 1957. Marine Algae of the Northeastern Coast of North America. University of Michigan Press, Ann Arbor. viii + 509 pp.
- Taylor, W. R. 1960. Marine Algae of the Eastern Tropical and Subtropical Coasts of the Americas. University of Michigan Press, Ann Arbor. xi + 870 pp.
- Troll, V. W. 1931. Botanische Mitteilungen aus den Tropen. III. *Dictyotopsis propagulifera*, eine neue Brackwasseralge ostindischer Mangrovesgebiete. Flora 125: 474–502.
- Tseng, C. K., X. Bangmei, X. Enzhan, Z. Derui, Z. Junfun, Z. Bailin, Z. Jinhua. 1984. Rhodophyta. In Common Seaweeds of China (Tseng C. K., ed.), pp. 43–165. Science Press, Beijing.
- Tsuda, R. T. 1991. Catalog of the marine benthic algae from the Ryukyu Islands, Japan. Galaxea 10: 35–64.
- Tsuda, R. T. & F. O. Wray. 1977. Bibliography of marine benthic algae in Micronesia. Micronesica 13: 85–120.
- Umezaki, I. 1961. The Marine Blue-Green Algae of Japan. Memoirs of the College of Agriculture, Kyoto University 83(8): 149 pp.
- Weber-van Bosse, A. 1921. Liste des algues du Siboga. II. Rhodophyceae. Première partie. Protofloridae, Nemalionales, Cryptonemiales. Siboga-Expeditie Monographie 59b: 187–31, pls. VI–VIII.

- Weber-van Bosse, A. 1923. Liste des algues du Siboga. III. Rhodophyceae. Seconde partie. Ceramiales. Siboga-Expeditie Monographie 59c: 311-392, pl. IX-X.
- Weber-van Bosse, A. 1928. Liste des agues du Siboga. IV. Rhodophyceae. Troisième partie. Gigartinales et Rhodymeniales et tableau de . . . Siboga-Expedite Monographie 59d: 393-533 pls. XI-XVI.
- Woelkerling, W. J. 1973. The morphology and systematics of the *Audouinella* complex (Acrochaetiaceae, Rhodophyta) in northeastern United States. *Rhodora* 75(804): 529-621.
- Womersley, H. B. S. 1971. The genus *Plocamium* (Rhodophyta) in southern Australia. *Trans. R. Soc. S. Aust.* 95(1): 9-27.
- Womersley, H. B. S. & A. Bailey. 1970. Marine algae of the Solomon Islands. *Roy. Soc. Lond. Phil. Trans. Ser. B. Biological Series* 259: 257-352.
- Wynne, M. J. & G. T. Kraft. 1985. *Hypoglossum caloglossoides* sp. nov. (Delesseriaceae, Rhodophyta) from Lord Howe Island, South Pacific. *Br. Phycol. J.* 20: 9-19.
- Yendo, K. 1920. Novae algae Japoniae. Decas I-III. *Bot. Mag. [Tokyo]*. 34: 1-12.

Received 7 May 1993.